


Arc Length

Alvin Lin

Calculus II: August 2016 - December 2016

Arc Length



Suppose we want to find the distance between $f(a)$ and $f(b)$ along the curve.

$$L = \int_a^b \sqrt{1 + f'(x)^2} dx$$

The arc length function is denoted:

$$S(x) = \int_a^x \sqrt{1 + f'(t)^2} dt$$

Example

$$f(x) = y = \ln(1 - x^2) \quad 0 \leq x \leq \frac{1}{2}$$

$$L = \int_a^b \sqrt{1 + f'(x)^2} dx$$

$$f'(x) = \frac{0 - 2x}{1 - x^2} = \frac{-2x}{1 - x^2}$$

$$\begin{aligned} 1 + f'(x)^2 &= 1 + \left(\frac{-2x}{1 - x^2}\right)^2 \\ &= 1 + \frac{4x^2}{(1 - x^2)^2} \\ &= \frac{(1 - x^2)^2 + 4x^2}{(1 - x^2)^2} \\ &= \frac{1 + x^4 - 2x^2 + 4x^2}{(1 - x^2)^2} \\ &= \frac{x^4 + 2x^2 + 1}{(1 - x^2)^2} \\ &= \frac{(1 + x^2)^2}{(1 - x^2)^2} \\ &= \left(\frac{1 + x^2}{1 - x^2}\right)^2 \end{aligned}$$

Therefore:

$$\begin{aligned} L &= \int_0^{\frac{1}{2}} \sqrt{1 + f'(x)^2} dx \\ &= \int_0^{\frac{1}{2}} \frac{1 + x^2}{1 - x^2} dx \\ &= \int_0^{\frac{1}{2}} \frac{2 - (1 - x^2)}{1 - x^2} dx \\ &= \int_0^{\frac{1}{2}} \frac{2}{1 - x^2} - \frac{1 - x^2}{1 - x^2} dx \\ &= \int_0^{\frac{1}{2}} \frac{2}{1 - x^2} - 1 dx \\ &= \int_0^{\frac{1}{2}} \frac{2}{(1 - x)(1 + x)} - 1 dx \end{aligned}$$

By partial fractions:

$$\int_0^{\frac{1}{2}} \frac{A}{1-x} + \frac{B}{1+x} - 1 dx$$

$$A = B = 1$$

$$\left[-A \ln |1-x| + B \ln |1+x| - x \right]_0^{\frac{1}{2}}$$

$$L = -\ln \left| \frac{1}{2} \right| + \ln \left| \frac{3}{2} \right| - \frac{1}{2}$$

You can find all my notes at <http://omgimanerd.tech/notes>. If you have any questions, comments, or concerns, please contact me at alvin@omgimanerd.tech