

Sequences

Alvin Lin

Calculus II: August 2016 - December 2016

Sequences

A list of numbers written in a definite order. Example:

$$\left\{ 1, 19, 11, 12, 19, 13, \dots, \frac{1}{11}, 19^2 \right\}$$

$$\left\{ 1, \frac{1}{2}, \frac{1}{3}, \frac{1}{4}, \dots \right\}$$

Define a mapping:

$$f : N \rightarrow R$$

$$f(n) = a_n$$

$$\left\{ a_1, a_2, a_3, a_4, \dots, a_n \right\}$$

a_n is the generic term where $a_n = \frac{1}{n}$.

Definition

A sequence $\{a_n\}$ has the limit L if we can make the terms as close to L as we want by choosing a sufficiently large n . If a_n has a limit L , we say that $\{a_n\}$ converges to L . We denote this by $a_n \rightarrow L$.

If there is no real number L such that $a_n \rightarrow L$, then $\{a_n\}$ diverges.

Example:

$$a_n = \frac{1}{n}$$

$$a_n \rightarrow 0$$

$$\lim_{n \rightarrow \infty} a_n = L$$

- a_n is increasing iff $a_n < a_{n+1} \quad \forall \quad n$.
- a_n is decreasing iff $a_n > a_{n+1} \quad \forall \quad n$.
- a_n is monotonic if it is either increase or decreasing.
- a_n is bounded from above iff $a_n \leq M \quad \forall \quad n$.
- a_n is bounded from below iff $a_n \geq M \quad \forall \quad n$.
- r^n is convergent is $-1 < r \leq 1$ and divergent for all other r 's.

Identities

- $\lim ca_n = c \lim a_n$
- $\lim a_n \pm b_n = \lim a_n + \lim b_n$
- $\lim \frac{a_n}{b_n} = \frac{\lim a_n}{\lim b_n}$
- $\lim a_n b_n = (\lim a_n)(\lim b_n)$
- $\lim ((a_n)^p) = (\lim a_n)^p \quad (p > 0, a_n > 0)$

Theorems

- If $\lim_{x \rightarrow \infty} f(x) = L$, then $a_n \rightarrow L$ where $a_n = f(n)$.
- If $\lim |a_n| = 0$, then $\lim a_n = 0$.
- Every bounded monotonic sequence is convergent. (Bounded from above and below). $|a_n| < M$.
- The Squeeze Theorem:

$$\begin{aligned}
 a_n &\leq b_n \leq c_n \quad n \geq n_o \\
 \lim_{n \rightarrow \infty} a_n &= \lim_{n \rightarrow \infty} c_n = L \\
 &\rightarrow \lim_{n \rightarrow \infty} b_n = L
 \end{aligned}$$

Practice Problem 17

$$a_n = 1 - (0.2)^n$$

$$\begin{aligned}\lim_{n \rightarrow \infty} a_n &= \lim_{n \rightarrow \infty} (1 - (0.2)^n) \\ &= 1 - \lim_{n \rightarrow \infty} (0.2)^n \\ &= 1 - 0 \\ &= 1\end{aligned}$$

$$\lim_{n \rightarrow \infty} r^n = \begin{cases} 0 & -1 < r < 1 \\ 1 & r = 1 \end{cases}$$

Any limit of this form converges when $-1 < r < 1$ otherwise it diverges.

Practice Problem 18

$$a_n = \frac{n^3}{n^3 + 1}$$

$$\begin{aligned}\lim_{n \rightarrow \infty} a_n &= \lim_{n \rightarrow \infty} \frac{n^3}{n^3 + 1} \\ &= \lim_{n \rightarrow \infty} \frac{n^3 \frac{1}{n^3}}{n^3 + 1 \frac{1}{n^3}} \\ &= \lim_{n \rightarrow \infty} \frac{1}{1 + \frac{1}{n^3}} \\ &= \frac{1}{1 + 0} \\ &= 1\end{aligned}$$

Practice Problem 21

$$a_n = e^{\frac{1}{n}}$$

$$\begin{aligned}\lim_{n \rightarrow \infty} a_n &= \lim_{n \rightarrow \infty} e^{\frac{1}{n}} \\ &= e^{\lim_{n \rightarrow \infty} \frac{1}{n}} \\ &= e^0 \\ &= 1\end{aligned}$$

Practice Problem 25

$$a_n = \frac{(-1)^n n}{n^2 + 1}$$

$$\begin{aligned}\lim_{n \rightarrow \infty} a_n &= \lim_{n \rightarrow \infty} \frac{(-1)^n n}{n^2 + 1} \\ \lim_{n \rightarrow \infty} |a_n| &= \lim_{n \rightarrow \infty} \frac{n}{n^2 + 1} \\ &= \lim_{n \rightarrow \infty} \frac{n}{n^2 + 1} \frac{\frac{1}{n^2}}{\frac{1}{n^2}} \\ &= \lim_{n \rightarrow \infty} \frac{\frac{1}{n}}{1 + \frac{1}{n^2}} \\ &= \frac{0}{1 + 0} \\ &= 0\end{aligned}$$

$$\lim_{n \rightarrow \infty} |a_n| = 0 \quad \therefore \quad \lim_{n \rightarrow \infty} a_n = 0$$

(Theorem)

Practice Problem 29

$$a_n = \frac{(2n - 1)!}{(2n + 1)!}$$

$$\begin{aligned}\lim_{n \rightarrow \infty} a_n &= \lim_{n \rightarrow \infty} \frac{(2n - 1)!}{(2n + 1)!} \\ &= \lim_{n \rightarrow \infty} \frac{(2n - 1)!}{(2n + 1)(2n)(2n - 1)!} \\ &= \lim_{n \rightarrow \infty} \frac{1}{(2n + 1)(2n)} \\ &= 0\end{aligned}$$

Practice Problem 36

$$a_n = \ln(n + 1) - \ln(n)$$

$$\begin{aligned}
\lim_{n \rightarrow \infty} a_n &= \lim_{n \rightarrow \infty} \ln(n+1) - \ln(n) \\
&= \lim_{n \rightarrow \infty} \ln\left(\frac{n+1}{n}\right) \\
&= \lim_{n \rightarrow \infty} \ln\left(1 + \frac{1}{n}\right) \\
&= \ln(1) \\
&= 0
\end{aligned}$$

Practice Problem 37

$$a_n = n \sin\left(\frac{1}{n}\right)$$

$$\begin{aligned}
\lim_{n \rightarrow \infty} a_n &= \lim_{x \rightarrow \infty} x \sin\left(\frac{1}{x}\right) \\
&= \lim_{x \rightarrow \infty} \frac{\sin\left(\frac{1}{x}\right)}{\frac{1}{x}} \\
&= \lim_{y \rightarrow 0} \frac{\sin(y)}{y} \\
&\quad \text{(L'Hopital's Rule)} \\
&= \lim_{y \rightarrow 0} \frac{\cos(y)}{1} \\
&= \lim_{t \rightarrow 0} \cos(y) \\
&= 1
\end{aligned}$$

Practice Problem 38

$$a_n = \sqrt[n]{2^{1+3n}}$$

$$\begin{aligned}
\lim_{n \rightarrow \infty} a_n &= \lim_{n \rightarrow \infty} 2^{\frac{1+3n}{n}} \\
&= \lim_{n \rightarrow \infty} 2^{\frac{1}{n}+3} \\
&= 2^{0+3} \\
&= 8
\end{aligned}$$

Practice Problem 39

$$a_n = \left(1 + \frac{2}{n}\right)^n$$
$$\lim_{n \rightarrow \infty} \left(1 + \frac{2}{n}\right)^n$$
$$y = y(x) = \left(1 + \frac{2}{x}\right)^x$$
$$\ln(y) = x \ln\left(1 + \frac{2}{x}\right)$$
$$\ln(y) = \frac{\ln\left(1 + \frac{2}{x}\right)}{\frac{1}{x}}$$

By L'Hopital's Rule:

$$\lim_{x \rightarrow \infty} \ln(y) = \lim_{x \rightarrow \infty} \frac{\ln\left(1 + \frac{2}{x}\right)}{\frac{1}{x}} = 2$$
$$\ln(y) = 2$$
$$y = e^2$$
$$\lim_{n \rightarrow \infty} a_n = e^2$$

Practice Problem 40

$$a_n = \frac{\sin(2n)}{1 + \sqrt{n}}$$
$$0 \leq a_n \leq \frac{1}{1 + \sqrt{n}}$$

By the Squeeze Theorem, since 0 and $\frac{1}{1+\sqrt{n}}$ converge to 0, a_n must converge to 0.

Practice Problem 66

$$a_n = n + \frac{1}{n}$$

Either $a_n \geq a_{n+1}$ or $a_{n+1} \geq a_n$ for all n.

$$f(x) = x + \frac{1}{x}$$
$$f'(x) = 1 + (-x^{-2}) = 1 - \frac{1}{x^2}$$
$$f'(x) > 0$$

We can conclude that a_n is increasing and has a lower bound of 0. Thus, a_n diverges.

Practice Problem 68

$$\begin{aligned}a_1 &= \sqrt{2} \\a_2 &= \sqrt{2 + \sqrt{2}} \\a_3 &= \sqrt{2 + \sqrt{2 + \sqrt{2}}} \\a_{n+1} &= \sqrt{2 + a_n} \\a_{n+2} &= \sqrt{2 + a_{n+1}}\end{aligned}$$

By the Monotonic Convergence Theorem:

$$\begin{aligned}\lim_{n \rightarrow \infty} a_n &= \lim_{n \rightarrow \infty} a_{n+1} = L \\ \lim_{n \rightarrow \infty} a_{n+1} &= \sqrt{2 + \lim_{n \rightarrow \infty} a_n} = \sqrt{2 + L} \\ L^2 &= 2 + L \\ L^2 - L - 2 &= 0 \\ (L - 2)(L + 1) &= 0 \\ L = 2 \quad L = -1\end{aligned}$$

We can eliminate -1 since we know the bounds of the sequence.

$$\lim_{n \rightarrow \infty} a_n = 2$$

You can find all my notes at <http://omgimanerd.tech/notes>. If you have any questions, comments, or concerns, please contact me at alvin@omgimanerd.tech