

Multivariable and Vector Calculus

Alvin Lin

August 2017 - December 2017

Cylindrical Coordinates

$$(x, y, z) \rightarrow (r, \theta, z)$$

Cylindrical coordinates describe a point by using polar coordinates and an elevation.

$$\begin{aligned}r &= \sqrt{x^2 + y^2} \\ \theta &= \tan^{-1}\left(\frac{y}{x}\right) \\ z &= z\end{aligned}$$

Theorem

$$\iiint_V f(x, y, z) \, dV = \iiint_C f(r \cos \theta, r \sin \theta, z) r \, dz \, d\theta \, dr$$

Spherical Coordinates

The distance between the origin and a point $P(x, y, z)$ is ρ . We can project this point onto the xy-plane to point $P'(x, y, 0)$. We can compute the polar angle θ between P' and the x-axis. If we fix the distance ρ and let all other variables free, we get a sphere. If we fix θ and ρ , the resulting set of points yields a semicircle. If we take the angle ϕ between P and the y-axis, we get a system of coordinates (ρ, θ, ϕ) described

by:

$$\begin{aligned}\rho &= \sqrt{x^2 + y^2 + z^2} \\ \theta &= \tan^{-1}\left(\frac{y}{x}\right) \\ \phi &= \cos^{-1}\left(\frac{z}{\sqrt{x^2 + y^2 + z^2}}\right) = \cos^{-1}\left(\frac{z}{\rho}\right)\end{aligned}$$

Converting the other way:

$$\begin{aligned}x &= \rho \sin \phi \cos \theta \\ y &= \rho \sin \phi \sin \theta \\ z &= \rho \cos \phi \\ \frac{d(x, y, z)}{d(\rho, \theta, \phi)} &= \rho^2 \sin \phi\end{aligned}$$

Example

$$\begin{aligned}(x, y, z) &\rightarrow (\rho, \theta, \phi) \\ P(2, 2, 3) &\equiv \left(\sqrt{17}, \frac{\pi}{4}, \cos^{-1}\left(\frac{3}{\sqrt{17}}\right)\right)\end{aligned}$$

$\rho = 1$: Sphere

$\phi = \frac{\pi}{4}$: Cone

$\phi = \frac{\pi}{2}$: xy-plane

$\phi = \frac{3\pi}{4}$: Cone

$\phi = \pi$: Negative z-axis

$\phi = 0$: Positive z-axis

Theorem

$$\begin{aligned}\iiint_V f(x, y, z) \, dV &= \iiint_C f(r \cos \theta, r \sin \theta, z) r \, dr \, d\theta \, dz \\ &= \iiint_S f(\rho \sin \phi \cos \theta, \rho \sin \phi \sin \theta, \rho \cos \phi) \rho^2 \sin \phi \, d\rho \, d\phi \, d\theta\end{aligned}$$

Average Value of a Function

$$A_v f = \frac{\iint_D f \, dA}{\iint_D 1 \, dA} = \frac{\iiint_V f \, dV}{\iiint_V 1 \, dV}$$

Center of Mass

$$\bar{x} = \frac{M_y}{M} = \frac{\iint_D f \cdot x \, dA}{\iint_D f \, dA}$$

Moment of Inertia

$$J_y = \iint_D f \cdot x^2 \, dA$$

Example

Find the center of mass of the solid bounded by the cone $r = 1, h = 1, f(x, y) = \sqrt{x^2 + y^2}$.

$$\begin{aligned}\bar{x} &= \bar{y} = 0 \\ \bar{z} &= \frac{M_{xy}}{\text{mass}} \\ &= \frac{\iiint_V \sqrt{x^2 + y^2} z \, dV}{\iiint_V \sqrt{x^2 + y^2} \, dV} \\ &= \frac{\int_0^{2\pi} \int_0^1 \int_r^1 r^2 z \, dz \, dr \, d\theta}{\int_0^{2\pi} \int_0^1 \int_r^1 r^2 \, dz \, dr \, d\theta}\end{aligned}$$

Note that this can also be represented by spherical coordinates:

$$= \frac{\int_0^{2\pi} \int_0^{\frac{\pi}{4}} \int_0^{\sec \phi} \rho^4 \sin^3 \phi \cos \phi \, d\rho \, d\phi \, d\omega}{\int_0^{2\pi} \int_0^{\frac{\pi}{4}} \int_0^{\sec \phi} \rho^3 \sin^3 \phi \, d\rho \, d\phi \, d\theta}$$

Example

Find the volume under $z = xy^2$ bounded by $z = 0$ and the triangle $(1,0), (2,0), (4,0)$.

$$\begin{aligned}V &= \iiint_V 1 \, dV \\ &= \int_0^1 \int_{1+y}^{4-2y} \int_0^{x^2 y} 1 \, dz \, dx \, dy\end{aligned}$$

You can find all my notes at <http://omgimanagerd.tech/notes>. If you have any questions, comments, or concerns, please contact me at alvin@omgimanagerd.tech