

Differential Equations

Alvin Lin

January 2018 - May 2018

Laplace Transforms

Let f be a function defined on $[0, \infty)$. The Laplace transform of f , $\mathcal{L}\{f(s)\}$, is a function $F(s)$, where s is the new variable. The domain of $\mathcal{L}\{f\}$, or $F(s)$, is all values of s for which the following converges:

$$\mathcal{L}\{f\} = \int_0^{\infty} e^{-st} f(t) dt$$

Recall the following:

$$\int_0^{\infty} g(t) dt = \lim_{N \rightarrow \infty} \int_0^N g(t) dt$$

If this limit exists, it converges. Otherwise, it diverges.

Example

Find $\mathcal{L}\{f(t)\}$ where $f(t) = 1$.

$$\begin{aligned} \mathcal{L}\{1\} &= \int_0^{\infty} e^{-st} dt \\ &= \lim_{N \rightarrow \infty} \int_0^{\infty} e^{-st} dt \\ &= \lim_{N \rightarrow \infty} \left[-\frac{1}{s} e^{-st} \right]_0^N \\ &= \lim_{N \rightarrow \infty} \left[-\frac{1}{e^{sN}} - \left(-\frac{1}{s} \right) \right] \\ &= \frac{1}{s}, \quad s > 0 \end{aligned}$$

Additional fact: $\mathcal{L}\{0\} = 0$.

Example

Find $\mathcal{L}\{e^{at}\}$ where a is real.

$$\begin{aligned}f(t) &= e^{at} \\ \mathcal{L}\{e^{at}\} &= \int_0^{\infty} e^{-st} e^{at} dt \\ &= \int_0^{\infty} e^{-(s-a)t} dt \\ &= \lim_{N \rightarrow \infty} \int_0^N e^{-(s-a)t} dt \\ &= \lim_{N \rightarrow \infty} \left[-\frac{1}{s-a} e^{-(s-a)t} \right]_0^N \\ &= \lim_{N \rightarrow \infty} \left[-\frac{1}{s-a} e^{-(s-a)N} - \left(-\frac{1}{s-a} \right) \right] \\ &= \lim_{N \rightarrow \infty} -\frac{1}{s-a} e^{-(s-a)N} + \frac{1}{s-a} \\ &= \frac{1}{s-a}, \quad s-a > 0 \\ &= \frac{1}{s-a}, \quad s > a \\ \mathcal{L}\{e^{2t}\} &= \frac{1}{s-2} \\ \mathcal{L}\{e^{-2t}\} &= \frac{1}{s+2}\end{aligned}$$

Example

Find $\mathcal{L}\{t\}$.

$$\begin{aligned}f(t) &= t \\ \mathcal{L}\{t\} &= \int_0^{\infty} te^{-st} dt \\ &= \lim_{N \rightarrow \infty} \int_0^N te^{-st} dt\end{aligned}$$

$$\begin{aligned}
&= \lim_{N \rightarrow \infty} \left[-\frac{1}{s} e^{-st} t \right]_0^N - \int_0^N -\frac{1}{s} e^{-st} dt \\
&= \lim_{N \rightarrow \infty} \left[-\frac{1}{s} e^{-st} t + \frac{1}{s} \left(-\frac{1}{s} e^{-st} \right) \right]_0^N = \lim_{N \rightarrow \infty} \left[\left(-\frac{1}{s} e^{-sN} N \right) + \left(-\frac{1}{s^2} e^{-sN} - \frac{1}{s^2} \right) \right] \\
&= \frac{1}{s^2}, \quad s > 0
\end{aligned}$$

As a general rule, for $f(t) = t^n$ where n is a non-negative integer:

$$\begin{aligned}
\mathcal{L}\{t^n\} &= \frac{n!}{s^{n+1}} \\
\mathcal{L}\{t^3\} &= \frac{3!}{s^{3+1}} = \frac{6}{s^4}
\end{aligned}$$

Additionally, for sines and cosines:

$$\begin{aligned}
\mathcal{L}\{\sin(bt)\} &= \frac{b}{s^2 + b^2}, \quad s > 0, b \neq 0 \\
\mathcal{L}\{\cos(bt)\} &= \frac{s}{s^2 + b^2}, \quad s > 0, b \neq 0 \\
\mathcal{L}\{\sin(\sqrt{2}t)\} &= \frac{\sqrt{2}}{s^2 + 2} \\
\mathcal{L}\{\cos(\sqrt{2}t)\} &= \frac{s}{s^2 + 2}
\end{aligned}$$

Property of Linearity

Given $f(t), g(t)$ and k a constant:

1. $\mathcal{L}\{f(t) \pm g(t)\} = \mathcal{L}\{f(t)\} \pm \mathcal{L}\{g(t)\}$
2. $\mathcal{L}\{kf(t)\} = k\mathcal{L}\{f(t)\}$


Example

$$\begin{aligned}
&\mathcal{L}\{7 + 5e^{-7t} - 7\sin(3t) + 7t^2 + 5\cos(7t)\} \\
&= \mathcal{L}\{7\} + \mathcal{L}\{5e^{-7t}\} - \mathcal{L}\{7\sin(3t)\} + \mathcal{L}\{7t^2\} + \mathcal{L}\{5\cos(7t)\} \\
&= 7\mathcal{L}\{1\} + 5\mathcal{L}\{e^{-7t}\} - 7\mathcal{L}\{\sin(3t)\} + 7\mathcal{L}\{t^2\} + 5\mathcal{L}\{\cos(7t)\} \\
&= \frac{7}{s} + \frac{5}{s+7} - \frac{(7)(3)}{s^2+9} + \frac{(7)(2!)}{s^3} + \frac{5s}{s^2+49}, \quad s > 0
\end{aligned}$$

Note that each term has a different condition for s , so we need to take the intersection of the conditions.

Existence of the Transform

Consider a piecewise continuous function f over $[a, b]$. f has a finite number of jump discontinuities.



We can extend over $[0, \infty)$. A function f is piecewise continuous over $[0, \infty)$ if f is piecewise continuous over $[0, N]$ for all $N > 0$.

$$\int_a^d f(t) dt = \int_a^b f(t) dt + \int_b^c f(t) dt + \int_c^d f(t) dt$$

Exponential Order α

A function f is of exponential order α if there exists positive constants M and T such that $|f(t)| \leq Me^{\alpha t}$ for all $t > T$. For example:

$$\begin{aligned} f(t) &= e^{2t} \sin(t) \\ |e^{2t} \sin(t)| &= e^{2t} |\sin(2t)| \leq Me^{2t} = e^{2t} \text{ for all } T \\ |\sin(2t)| &\leq 1 \end{aligned}$$

$f(t)$ is of exponential order $\alpha = 2$.

Conditions for Existence of the Laplace Transform

1. If $f(t)$ is piecewise continuous and of exponential order α , then the Laplace transform $\mathcal{L}\{f(t)\}$ exists for $s > \alpha$.
2. We need to know that

$$\mathcal{L}\{f(t)\} = \int_0^{\infty} e^{-st} f(t) dt$$

converges for $s > a$. We can rewrite it as

$$\int_0^\infty e^{-st} f(t) dt = \int_0^T e^{-st} f(t) dt + \int_T^\infty e^{-st} f(t) dt$$

where we have $|f(t)| \leq Me^{\alpha t}$ for all $t > T$. By continuity, $\int_0^T e^{-st} f(t) dt$ exists, so we only need to consider $\int_T^\infty e^{-st} f(t) dt$.

$$\begin{aligned} \int_T^\infty e^{-st} f(t) dt &= \lim_{N \rightarrow \infty} \int_T^N e^{-st} f(t) dt \\ &\leq \lim_{N \rightarrow \infty} \int_T^N Me^{-st} e^{\alpha t} dt \\ &= \lim_{N \rightarrow \infty} M \int_T^N e^{-(s-\alpha)t} dt \\ &= \lim_{N \rightarrow \infty} M \left[-\frac{1}{s-\alpha} e^{-(s-\alpha)t} \right]_T^N \\ &= \lim_{N \rightarrow \infty} M \left[-\frac{1}{s-\alpha} e^{-(s-\alpha)N} - \left(-\frac{1}{s-\alpha} e^{-(s-\alpha)T} \right) \right] \\ &= \frac{M}{s-\alpha} e^{-(s-\alpha)T} < \infty \end{aligned}$$

Since M and T are positive constants, this integral converges by the comparison test for improper integrals.

Properties of the Transform

- Translation in s : If the Laplace transform $\mathcal{L}\{f(t)\} = F(s)$ exists, then $\mathcal{L}\{e^{\alpha t} f(t)\} = F(s - a)$. $e^{\alpha t}$ indicates a translation in the s domain, where we replace s by $s - a$ in the transform of $f(t)$. For example:

$$\begin{aligned} \mathcal{L}\{e^{2t} \sin(2t)\} & \quad \alpha = 2 \\ \mathcal{L}\{\sin(2t)\} &= \frac{2}{s^2 + 4} \\ \mathcal{L}\{e^{2t} \sin(2t)\} &= \frac{2}{(s - 2)^2 + 4} \end{aligned}$$

Another example:

$$\begin{aligned}\mathcal{L}\{e^{-2t} \cos(2t)\} & \quad \alpha = -2 \\ \mathcal{L}\{\cos(2t)\} & = \frac{s}{s^2 + 4} \\ \mathcal{L}\{e^{-2t} \cos(2t)\} & = \frac{s + 2}{(s + 2)^2 + 4}\end{aligned}$$

The Laplace Transform of the Derivative

Let f be continuous on $[0, \infty)$ and f' be piecewise continuous on $[0, \infty)$ and both are of exponential order α .

$$\begin{aligned}\mathcal{L}\{f'(t)\} & = \int_0^\infty e^{-st} f'(t) dt \\ & = \lim_{N \rightarrow \infty} \int_0^N e^{-st} f'(t) dt \\ u = e^{-st} & \quad dv = f'(t) dt \\ du = -se^{-st} dt & \quad v = f(t) \\ & = \lim_{N \rightarrow \infty} \left[e^{-st} f(t) \right]_0^N + \lim_{N \rightarrow \infty} s \int_0^N e^{-st} f(t) dt \\ & = -f(0) + sF(s) \\ & = sF(s) - f(0)\end{aligned}$$

In summary:

$$\begin{aligned}\mathcal{L}\{y'\} & = sY(s) - y(0) \\ \mathcal{L}\{y''\} & = s^2Y(s) - sy(0) - y'(0)\end{aligned}$$

Derivatives of the Laplace Transform

Let $F(S) = \mathcal{L}\{f(t)\}$. Then for $s > a$:

$$\mathcal{L}\{t^n f(t)\} = (-1)^n \frac{d^n F}{ds^n}$$

For example:

$$\begin{aligned}
 \mathcal{L}\{t \sin(t)\} \quad n = 1 \quad f(t) = \sin(t) \\
 \mathcal{L}\{\sin(t)\} &= \frac{1}{s^2 + 1} = F(s) \\
 \frac{dF}{ds} &= \frac{d}{ds} \left[\frac{1}{s^2 + 1} \right] \\
 &= \frac{(s^2 + 1)(0) - 1(2s)}{(s^2 + 1)^2} \\
 &= \frac{-2s}{(s^2 + 1)^2} \\
 \mathcal{L}\{t \sin(t)\} &= (-1)^1 \frac{-2s}{(s^2 + 1)^2} = \frac{2s}{(s^2 + 1)^2}
 \end{aligned}$$

Another example:

$$\begin{aligned}
 \mathcal{L}\{te^{-2t} \sin(3t)\} \quad n = 1 \quad f(t) = e^{-2t} \sin(3t) \\
 F(s) = \mathcal{L}\{e^{-2t} \sin(3t)\} \\
 \mathcal{L}\{\sin(3t)\} &= \frac{3}{s^2 + 9} \\
 \mathcal{L}\{e^{-2t} \sin(3t)\} &= \frac{3}{(s + 2)^2 + 9} = F(s) \\
 \frac{d}{ds} F(s) &= \frac{0 - (3(2(s + 2)))}{((s + 2)^2 + 9)^2} \\
 &= \frac{-6(s + 2)}{((s + 2)^2 + 9)^2} \\
 \mathcal{L}\{te^{-2t} \sin(3t)\} &= (-1)^1 \frac{-6(s + 2)}{((s + 2)^2 + 9)^2} \\
 &= \frac{6(s + 2)}{((s + 2)^2 + 9)^2}
 \end{aligned}$$

Inverse Laplace Transforms

The Laplace transform maps a function $f(t)$ to a function $F(s) = \mathcal{L}\{f(t)\}$. The transform has an inverse denoted by:

$$\mathcal{L}^{-1}\{f(t)\} = f(t)$$

Consider the following initial value problem:

$$y'' - y = -t \quad y(0) = 0 \quad y'(0) = 1$$

If we apply the Laplace transform to the entire equation:

$$\begin{aligned}\mathcal{L}\{y'' - y\} &= \mathcal{L}\{-t\} \\ \mathcal{L}\{y''\} - \mathcal{L}\{y\} &= -\mathcal{L}\{t\} \\ s^2Y(s) - (s)y(0) - y'(0) - Y(s) &= -\frac{1}{s^2} \\ (s^2 - 1)Y(s) &= -\frac{1}{s^2} + 1 \\ &= \frac{-1 + s^2}{s^2} \\ Y(s) &= \frac{1}{s^2} \\ \mathcal{L}^{-1}\{Y(s)\} &= \mathcal{L}^{-1}\left\{\frac{1}{s^2}\right\} \\ t &= y(t)\end{aligned}$$

Given $F(s)$, if there exists a function $f(t)$ that is continuous on $[0, \infty)$ where $\mathcal{L}\{f(t)\} = F(s)$, $f(t)$ is the inverse of $F(s)$. The inverse Laplace transform is a linear operator. Given $F_1(s), F_2(s)$ and k a constant, the properties of linearity apply:

1. $\mathcal{L}^{-1}\{F_1(s) \pm F_2(s)\} = \mathcal{L}^{-1}\{F_1(s)\} \pm \mathcal{L}^{-1}\{F_2(s)\}$
2. $\mathcal{L}^{-1}\{kF_1(s)\} \pm = k\mathcal{L}^{-1}\{F_1\}$

Example

$$\begin{aligned}\mathcal{L}^{-1}\left\{\frac{3}{s^2 + 9}\right\} &= \mathcal{L}^{-1}\left\{\frac{3}{s^2 + 3^2}\right\} \\ &= \sin(3t) \\ \mathcal{L}^{-1}\left\{\frac{s - 2}{(s - 2)^2 + 4}\right\} &= e^{2t} \cos(2t) \\ \mathcal{L}^{-1}\left\{\frac{1}{s - 6}\right\} &= e^{6t}\end{aligned}$$

Review of Partial Fraction Decomposition: Case 1

Suppose we have $f(x) = \frac{P(x)}{Q(x)}$ and the degree of P is less than the degree of Q .

$$\frac{P(x)}{Q(x)} = \frac{P}{(a_1x + b_1)(a_2x + b_2) \cdots (a_nx + b_n)}$$

We have n distinct linear factors that are irreducible, which we can rewrite as:

$$\frac{P(x)}{Q(x)} = \frac{A_1}{a_1x + b_1} + \frac{A_2}{a_2x + b_2} + \cdots + \frac{A_n}{a_nx + b_n}$$

where all A_i 's are constants. For example:

$$\begin{aligned} \frac{x + 4}{x^2 + 3x + 2} &= \frac{x + 4}{(x + 1)(x + 2)} \\ &= \frac{A}{x + 1} + \frac{B}{x + 2} = \frac{A(x + 2) + B(x + 1)}{(x + 1)(x + 2)} \\ x + 4 &= A(x + 2) + B(x + 1) \\ \text{Let : } x &= -2 \\ -2 + 4 &= -B \quad B = -2 \\ \text{Let : } x &= -1 \\ A &= 3 \end{aligned}$$

This can be applied to inverse Laplace transforms:

$$\begin{aligned} \mathcal{L}^{-1}\left\{\frac{3s + 2}{s^2 + s - 2}\right\} \\ \frac{3s + 2}{s^2 + s - 2} &= \frac{3s + 2}{(s + 2)(s - 1)} \\ &= \frac{A}{s + 2} + \frac{B}{s - 1} \\ &= \frac{A(s - 1) + B(s + 2)}{(s + 2)(s - 1)} \\ 3s + 2 &= A(s - 1) + B(s + 2) \\ \text{Let : } s &= 1 \quad 5 = 3B \quad B = \frac{5}{3} \\ \text{Let : } s &= -2 \quad A = \frac{4}{3} \end{aligned}$$

$$\begin{aligned}
\mathcal{L}^{-1}\left\{\frac{3s+2}{s^2+s-2}\right\} &= \mathcal{L}^{-1}\left\{\frac{4}{3}\frac{1}{s-1} + \mathcal{L}^{-1}\left\{\frac{5}{3}\frac{1}{s+2}\right\}\right\} \\
&= \frac{4}{3}\mathcal{L}^{-1}\left\{\frac{1}{s-1}\right\} + \frac{5}{3}\mathcal{L}^{-1}\left\{\frac{1}{s+2}\right\} \\
&= \frac{4}{3}e^t + \frac{5}{3}e^{-2t}
\end{aligned}$$

Review of Partial Fraction Decomposition: Case 2

In this case, we can have n repeated linear factors:

$$f(x) = \frac{P(x)}{Q(x)} = \frac{P(x)}{(a_1x + b_1)^n} = \frac{A_1}{a_1x + b_1} + \frac{A_2}{(a_1x + b_1)^2} + \cdots + \frac{A_n}{(a_1x + b_1)^n}$$

For example:

$$\mathcal{L}^{-1}\left\{\frac{s^2 + 9s + 2}{(s-2)^2(s+3)}\right\}$$

$$\frac{A}{s+3} + \frac{B}{s-1} + \frac{C}{(s-1)^2} = \frac{A(s-1)^2 + B(s+3)(s-1) + C(s+3)}{(s-1)^2(s+3)}$$

$$s^2 + 9s + 2 = A(s-1)^2 + B(s+3)(s-1) + C(s+3)$$

$$\text{Let : } s = 1$$

$$1 + 9 + 2 = 4C \quad 4C = 12 \quad C = 3$$

$$\text{Let : } s = -3$$

$$9 - 27 + 2 = 16A \quad A = -1$$

$$\text{Let : } s = 0$$

$$2 = -1 - 3B + 3C \quad B = 2$$

$$\mathcal{L}^{-1}\left\{\frac{s^2 + 9s + 2}{(s-2)^2(s+3)}\right\} = \mathcal{L}^{-1}\left\{\frac{-1}{s+3} + \frac{2}{s-1} + \frac{3}{(s-1)^2}\right\}$$

$$= -\mathcal{L}^{-1}\left\{\frac{1}{s+3}\right\} + 2\mathcal{L}^{-1}\left\{\frac{1}{s-1}\right\} + 3\mathcal{L}^{-1}\left\{\frac{1}{(s-1)^2}\right\}$$

$$= -e^{-3t} + 2e^t + 3te^t$$

Review of Partial Fraction Decomposition: Case 3

In this case, we have n irreducible distinct quadratics:

$$\begin{aligned}\frac{P(x)}{Q(x)} &= \frac{P(x)}{(a_1x^2 + b_1x + c) \cdots (a_nx^2 + b_nx + c_n)} \\ &= \frac{A_1x + B_1}{a_1x^2 + b_1x + c_1} + \cdots + \frac{A_nx + B_n}{a_nx^2 + b_nx + c_n}\end{aligned}$$

For example:

$$\mathcal{L}^{-1}\left\{\frac{7s - 8}{(s^2 + 4)(s + 1)(s - 2)}\right\}$$

$$\begin{aligned}\frac{7s - 8}{(s^2 + 4)(s + 1)(s - 2)} &= \frac{A}{s - 1} + \frac{B}{s - 2} + \frac{Cs + D}{s^2 + 4} \\ 7s - 8 &= A(s - 2)(s^2 + 4) + B(s + 1)(s^2 + 4) + (Cs + D)(s + 1)(s - 2)\end{aligned}$$

$$\text{Let : } s = 2 \quad B = \frac{1}{4}$$

$$\text{Let : } s = -1 \quad A = 1$$

$$\text{Let : } s = 0 \quad D = \frac{1}{2}$$

$$\text{Let : } s = 1 \quad C = -\frac{5}{4}$$

$$\begin{aligned}\mathcal{L}^{-1}\left\{\frac{7s - 8}{(s^2 + 4)(s + 1)(s - 2)}\right\} &= \mathcal{L}^{-1}\left\{\frac{1}{s + 1}\right\} + \mathcal{L}^{-1}\left\{\frac{1}{4} \frac{1}{s - 2}\right\} + \mathcal{L}^{-1}\left\{\frac{-\frac{5}{4}s + \frac{1}{2}}{s^2 + 4}\right\} \\ &= \mathcal{L}^{-1}\left\{\frac{1}{s + 1}\right\} + \frac{1}{4}\mathcal{L}^{-1}\left\{\frac{1}{s - 2}\right\} - \frac{5}{4}\mathcal{L}^{-1}\left\{\frac{s}{s^2 + 4}\right\} + \frac{1}{2}\mathcal{L}^{-1}\left\{\frac{1}{s^2 + 4}\right\} \\ &= e^{-t} + \frac{1}{4}e^{2t} - \frac{5}{4}\cos(2t) + \frac{1}{4}\sin(2t)\end{aligned}$$

Solving Differential Equations with Laplace Transforms

We can use the Laplace transform to solve certain initial value problems. The general strategy is to take the Laplace transform of both sides of the equation, simplify, and then take the inverse Laplace transform. Given the following initial value problem:

$$y'' - y' - 2y = 0 \quad y(0) = -2 \quad y'(0) = 5$$

First we take the Laplace transform of both sides.

$$\begin{aligned}\mathcal{L}\{y'' - y' - 2y\} &= \mathcal{L}\{0\} \\ \mathcal{L}\{y''\} - \mathcal{L}\{y'\} - 2\mathcal{L}\{y\} &= \mathcal{L}\{0\}\end{aligned}$$

We can solve for the Laplace transforms and isolate $Y(s)$, which is the Laplace transform of our solution $y(t)$.

$$\begin{aligned}\left[s^2Y(s) - sy(0) - y'(0)\right] - \left[sY(s) - y(0)\right] - 2(Y(s)) &= 0 \\ s^2Y(s) + 2s - 5 - sY(s) - 2 - 2Y(s) &= 0 \\ s^2Y(s) - sY(s) - 2Y(s) &= 7 - 2s \\ Y(s)(s^2 - s - 2) &= 7 - 2s \\ Y(s) &= \frac{7 - 2s}{s^2 - s - 2}\end{aligned}$$

Now we simply take the inverse Laplace transform of $Y(s)$ using partial fraction decomposition to get our solution $y(t)$.

$$\begin{aligned}y(t) &= \mathcal{L}^{-1}\{Y(s)\} \\ \frac{A}{s-2} + \frac{B}{s+1} &= \frac{7-2s}{(s-2)(s+1)} \\ A(s+1) + B(s-2) &= 7-2s \\ \text{Let : } s = -1 \quad B &= -3 \\ \text{Let : } s = 2 \quad A &= 1 \\ y(t) &= \mathcal{L}^{-1}\left\{\frac{1}{s-2} - \frac{3}{s+1}\right\} \\ &= \mathcal{L}^{-1}\left\{\frac{1}{s-2}\right\} - 3\mathcal{L}^{-1}\left\{\frac{1}{s+1}\right\} \\ &= e^{2t} - 3e^{-t}\end{aligned}$$

This strategy can be used to solve other differential equations of similar form.

Transforms of Discontinuous Functions

First, let's introduce the idea of the unit step function:

$$u(t) = \begin{cases} 0, & t < 0 \\ 1, & t \geq 0 \end{cases}$$

For $a > 0$:

$$u(t - a) = \begin{cases} 0, & t < a \\ 1 & t \geq a \end{cases}$$

Suppose we have the function:

$$f(t) = \begin{cases} -1, & 0 \leq t < 3 \\ 1, & 3 < t < 5 \\ 0, & t \geq 5 \end{cases}$$

We can represent this function using the unit step function. The first jump discontinuity occurs at $t = 3$, which can be associated with the following unit step function:

$$u(t - 3) = \begin{cases} 0, & t < 3 \\ 1, & t \geq 3 \end{cases}$$

The jump discontinuity at $t = 5$ can be associated with the following:

$$u(t - 5) = \begin{cases} 0, & t < 5 \\ 1, & t \geq 5 \end{cases}$$

Using this, we can rewrite our discontinuous function as the following:

$$f(t) = -1 + 2u(t - 3) - u(t - 5)$$

At the boundary on $t = 3$, the function increases by a value of 2, which becomes the coefficient for $u(t - 3)$. At the boundary on $t = 5$, the function decreases by a value of 1, which becomes the coefficient for $u(t - 5)$.

Example

Let's consider another example. Suppose now we have the function:

$$f(t) = \begin{cases} 1, & 0 < t < 2 \\ t, & 2 < t < 3 \\ -1, & t \geq 3 \end{cases}$$

We can represent this function using the unit step function as follows:

$$f(t) = 1 + (t - 1)u(t - 2) - (t + 1)u(t - 3)$$

Laplace Transforms of the Unit Step

By definition:

$$\mathcal{L}\{u(t-a)\} = \int_0^{\infty} e^{-st} u(t-a) dt$$

Since this is 0 for $t < a$ by the definition of the unit step function, we can rewrite this as follows:

$$\begin{aligned} \int_0^{\infty} e^{-st} u(t-a) dt &= \int_a^{\infty} e^{-st} dt \\ &= \lim_{N \rightarrow \infty} \int_a^N e^{-st} dt \\ &= \lim_{N \rightarrow \infty} \left[-\frac{1}{s} e^{-st} \right]_a^N \\ &= \lim_{N \rightarrow \infty} \left[-\frac{1}{s} e^{-sN} - \left(-\frac{1}{s} e^{-sa}\right) \right] \\ &= \frac{e^{-as}}{s} \end{aligned}$$

For a function $y = f(t-a)u(t-a)$ translated in the t domain:

$$\mathcal{L}\{f(t-a)u(t-a)\} = e^{-as}F(s) = e^{-as}\mathcal{L}\{f(t)\}$$

Example

$$\begin{aligned} \mathcal{L}\{4u(t-2)\} &= e^{-2s}\mathcal{L}\{4\} \\ &= e^{-2s}\frac{4}{s} \\ \mathcal{L}\{\sin(t-\pi)u(t-\pi)\} &= e^{-\pi s}\mathcal{L}\{\sin(t)\} \\ &= e^{-\pi s}\frac{1}{s^2+1} \end{aligned}$$

The same logic can be applied for inverse transforms:

$$\begin{aligned} \mathcal{L}^{-1}\left\{e^{-\pi s}\frac{1}{(s-1)^2+1} + e^{-2\pi s}\frac{1}{s^2}\right\} &= \mathcal{L}^{-1}\left\{e^{-\pi s}\frac{1}{(s-1)^2+1}\right\} + \mathcal{L}^{-1}\left\{e^{-2\pi s}\frac{1}{s^2}\right\} \\ &= e^{t-\pi} \sin(t-\pi)u(t-\pi) + (t-2\pi)u(t-2\pi) \end{aligned}$$

Example

We can use these techniques to solve differential equations as well. Suppose we have:

$$\begin{aligned}y'' + y &= f(t) \\f(t) &= \begin{cases} 0, & 0 < t < 2 \\ 1, & 2 < t < 4 \\ 0, & 4 < t \end{cases} \\y(0) &= 1 \\y'(0) &= 0\end{aligned}$$

We can rewrite $f(t)$ using the unit step function.

$$f(t) = u(t - 2) - u(t - 4)$$

Now we can use the method of Laplace transforms to solve the differential equation.

$$\begin{aligned}y'' + y &= u(t - 2) - u(t - 4) \\ \mathcal{L}\{y'' + y\} &= \mathcal{L}\{u(t - 2)\} - \mathcal{L}\{u(t - 4)\} \\ \mathcal{L}\{y''\} + \mathcal{L}\{y\} &= \mathcal{L}\{u(t - 2)\} - \mathcal{L}\{u(t - 4)\} \\ \left[s^2 Y(s) - sy(0) - y'(0) \right] + Y(s) &= \frac{e^{-2s}}{s} - \frac{e^{-4s}}{s} \\ Y(s)(s^2 + 1) - s &= \frac{e^{-2s}}{s} - \frac{e^{-4s}}{s} \\ Y(s) &= \frac{s}{s^2 + 1} + \frac{e^{-2s}}{s(s^2 + 1)} - \frac{e^{-4s}}{s(s^2 + 1)} \\ y(t) &= \mathcal{L}^{-1}\{Y(s)\} \\ &= \mathcal{L}^{-1}\left\{\frac{s}{s^2 + 1}\right\} + \mathcal{L}^{-1}\left\{e^{-2s} \frac{1}{s(s^2 + 1)}\right\} - \mathcal{L}^{-1}\left\{e^{-4s} \frac{1}{s(s^2 + 1)}\right\}\end{aligned}$$

We will need to use partial fraction decomposition to solve these inverse transforms.

$$\begin{aligned}\frac{1}{s(s^2 + 1)} &= \frac{A}{s} + \frac{Bs + C}{s^2 + 1} \\ &= \frac{A(s^2 + 1) + (Bs + C)s}{s(s^2 + 1)}\end{aligned}$$

Let : $s = 0$ $A = 1$

$B = -1$

$C = 0$

$$\begin{aligned}y(t) &= \mathcal{L}^{-1}\left\{\frac{s}{s^2 + 1}\right\} + \mathcal{L}^{-1}\left\{e^{-2s}\left(\frac{1}{s} - \frac{s}{s^2 + 1}\right)\right\} - \mathcal{L}^{-1}\left\{e^{-4s}\left(\frac{1}{s} - \frac{s}{s^2 + 1}\right)\right\} \\ &= \cos(t) + u(t - 2) - \cos(t - 2)u(t - 2) - u(t - 4) + \cos(t - 4)u(t - 4)\end{aligned}$$

Example

$$y' - 4y = \sin(t)u(t - 2\pi) \quad y(0) = 1$$

$$y' - 4y = \sin(t - 2\pi)u(t - 2\pi)$$

$$\mathcal{L}\{y' - 4y\} = \mathcal{L}\{\sin(t - 2\pi)u(t - 2\pi)\}$$

$$\mathcal{L}\{y'\} - 4\mathcal{L}\{y\} = \mathcal{L}\{\sin(t - 2\pi)u(t - 2\pi)\}$$

$$sY(s) - 1 - 4Y(s) = e^{-2\pi s} \frac{1}{s^2 + 1}$$

$$(s - 4)Y(s) = 1 + e^{-2\pi s} \frac{1}{s^2 + 1}$$

$$Y(s) = \frac{1}{s - 4} + e^{-2\pi s} \frac{1}{(s - 4)(s^2 + 1)}$$

$$\frac{1}{(s - 4)(s^2 + 1)} = \frac{A}{s - 4} + \frac{Bs + C}{s^2 + 1}$$

$$1 = A(s^2 + 1) + (Bs + C)(s - 4)$$

$$A = -\frac{1}{17} \quad B = -\frac{4}{17} \quad C = \frac{1}{17}$$

$$\begin{aligned}
Y(s) &= \frac{1}{s-4} + e^{-2\pi s} \left[-\frac{1}{17} \frac{1}{s-4} - \frac{4}{17} \frac{s}{s^2+1} + \frac{1}{17} \frac{1}{s^2+1} \right] \\
y(t) &= \mathcal{L}^{-1}\{Y(s)\} \\
&= e^{4t} - \frac{1}{17} e^{4(t-2\pi)} u(t-2\pi) - \frac{4}{17} \cos(t-2\pi) u(t-2\pi) + \frac{1}{17} \sin(t-2\pi) u(t-2\pi)
\end{aligned}$$

In general, for transformations in the t domain, the following Laplace transform rules apply:

$$\begin{aligned}
\mathcal{L}\{u(t-a)\} &= \frac{e^{-as}}{s}, \quad a \geq 0 \\
\mathcal{L}\{f(t-a)u(t-a)\} &= e^{-as}F(s), \quad a > 0 \\
\mathcal{L}\{g(t)u(t-a)\} &= e^{-as}\mathcal{L}\{g(t+a)\}, \quad a > 0
\end{aligned}$$

Impulses and the Dirac Delta Function

The Dirac Delta function is defined as:

$$\delta(t-a) = \begin{cases} \infty, & t = a \\ 0, & \text{otherwise} \end{cases}$$

Property of $\delta(t-a)$:

$$\int_0^{\infty} f(t)\delta(t-a) dt = f(a)$$

It “shifts” out one value of f , out of all possible values of f , which is $f(a)$.

$$\mathcal{L}\{\delta(t-a)\} = \int_0^{\infty} e^{-st}\delta(t-a) dt = e^{-as}$$

Additionally:

$$\int_0^{\infty} \delta(t-a) dt = 1$$

Example

Consider the governing equation of the motion of a mass on a spring moving where damping is negligible. At $t = 2\pi$, the mass is given a sharp blow given by $4\delta(t - 2\pi)$. The mass is released from rest 1 unit below the equilibrium position (positive

direction) with no initial velocity. Find the equation of motion given that the mass is 1kg and the spring constant is 1 Newton per meter.

$$\begin{aligned}
 y'' + y &= 4\delta(t - 2\pi) \\
 \mathcal{L}\{y'' + y\} &= \mathcal{L}\{4\delta(t - 2\pi)\} \\
 \mathcal{L}\{y''\} + \mathcal{L}\{y\} &= 4\mathcal{L}\{\delta(t - 2\pi)\} \\
 s^2Y(s) - sy(0) - y'(0) + Y(s) &= 4e^{-2\pi s} \\
 s^2Y(s) - s - 0 + Y(s) &= 4e^{-2\pi s} \\
 Y(s)(s^2 + 1) &= s + 4e^{-2\pi s} \\
 Y(s) &= \frac{s}{s^2 + 1} + \frac{4e^{-2\pi s}}{s^2 + 1} \\
 y(t) &= \mathcal{L}^{-1}\{Y(s)\} \\
 \mathcal{L}^{-1}\left\{\frac{s}{s^2 + 1}\right\} &= \cos(t) \\
 4\mathcal{L}^{-1}\left\{\frac{e^{-2\pi s}}{s^2 + 1}\right\} &= 4\mathcal{L}^{-1}\{e^{-as}F(s)\} \\
 &= 4f(t - a)u(t - a) \\
 a &= 2\pi \\
 F(s) &= \frac{1}{s^2 + 1} \\
 f(t) &= \sin(t) \\
 4f(t - a)u(t - a) &= 4\sin(t - 2\pi)u(t - 2\pi) \\
 &= 4\sin(t)u(t - 2\pi) \\
 y(t) &= \cos(t) + 4\sin(t)u(t - 2\pi) \\
 &= \begin{cases} \cos(t), & 0 \leq t \leq 2\pi \\ \cos(t) + 4\sin(t), & t > 2\pi \end{cases}
 \end{aligned}$$


Convolution

Let f and g be piecewise continuous functions over $[0, \infty)$. The convolution of f and g is defined as:

$$f * g = \int_0^t f(t - v)g(v) \, dv$$

We call this the convolution integral. This operation is commutative, in other words:

$$g * f = f * g$$



This effect is known as smearing.

Example

Find:

$$\begin{aligned}
 e^{3t} * e^{7t} &= \int_0^t e^{3(t-v)} e^{7v} dt \\
 &= \int_0^t e^{3t} e^{-3v} e^{7v} dv \\
 &= e^{3t} \int_0^t e^{4v} dv \\
 &= e^{3t} \left[\frac{1}{4} e^{4v} \right]_0^t \\
 &= e^{3t} \left[\frac{1}{4} (e^{4t} - e^0) \right] \\
 &= \frac{e^{7t} - e^{3t}}{4}
 \end{aligned}$$

Example

Find:

$$\begin{aligned}f(t) &= e^t & g(t) &= e^{-2t} \\f * g &= \int_0^t e^{t-v} e^{-2v} \, dv \\&= \int_0^t e^t e^{-v} e^{-2v} \, dv \\&= e^t \int_0^t e^{-3v} \, dv \\&= e^t \left[-\frac{1}{3} e^{-3v} \right]_0^t \\&= \frac{e^t - e^{-2t}}{3}\end{aligned}$$

The Convolution Theorem

$$\mathcal{L}\{f * g\} = F(s)G(s)$$

Examples:

$$\mathcal{L}\{e^{2t} * \sin(t)\} = \frac{1}{s-2} \frac{1}{s^2+1}$$

$$\begin{aligned}\mathcal{L}^{-1}\left\{\frac{1}{s+1} \frac{1}{s+2}\right\} &= e^{-t} * e^{-2t} \\&= \int_0^t e^{-(t-v)} e^{-2v} \, dv \\&= \int_0^t e^{-t} e^v e^{-2v} \, dv \\&= e^{-t} \int_0^t e^{-v} \, dv \\&= e^{-t} \left[-e^{-v} \right]_0^t \\&= e^{-t} - e^{-2t}\end{aligned}$$

Example

Solve the following integral-differential equation using the method of Laplace transforms:

$$y' + 6y + 9 \int_0^t y(v) \, dv = 1 \quad y(0) = 0$$

$$\begin{aligned} \int_0^t y(v) \, dv &= \int_0^t 1y(v) \, dv \\ &= 1 * y(t) \end{aligned}$$

$$y' + 6y + 9(1 * y(t)) = 1$$

$$\mathcal{L}\{y' + 6y + 9(1 * y(t))\} = \mathcal{L}\{1\}$$

$$sY(s) + 6Y(s) + 9\left[\frac{1}{s}Y(s)\right] = \frac{1}{s}$$

$$Y(s)\left(s + 6 + \frac{9}{s}\right) = \frac{1}{s}$$

$$Y(s)\left(\frac{s^2 + 6s + 9}{s}\right) = \frac{1}{s}$$

$$Y(s) = \frac{1}{(s + 3)^2}$$

$$\begin{aligned} y(t) &= \mathcal{L}^{-1}\left\{\frac{1}{(s + 3)^2}\right\} \\ &= e^{-3t}t \end{aligned}$$

Example

Solve the following integral equation.

$$y(t) = \cos(t) + \int_0^t e^{-v} y(t-v) \, dv$$

$$\int_0^t e^{-v} y(t-v) \, dv = e^{-t} * y(t)$$

$$y(t) = \cos(t) + e^t * y(t)$$

$$\mathcal{L}\{y(t)\} = \mathcal{L}\{\cos(t) + e^t * y(t)\}$$

$$Y(s) = \frac{s}{s^2 + 1} + \frac{1}{s + 1} Y(s)$$

$$Y(s) - \frac{1}{s + 1} Y(s) = \frac{s}{s^2 + 1}$$

$$\left(1 - \frac{1}{s + 1}\right) Y(s) = \frac{s}{s^2 + 1}$$

$$\left(\frac{s + 1 - 1}{s + 1}\right) Y(s) = \frac{s}{s^2 + 1}$$

$$\frac{s}{s + 1} Y(s) = \frac{s}{s^2 + 1}$$

$$Y(s) = \frac{s}{s^2 + 1} \frac{s + 1}{s}$$

$$= \frac{s + 1}{s^2 + 1}$$

$$= \frac{s}{s^2 + 1} + \frac{1}{s^2 + 1}$$

$$y(t) = \mathcal{L}^{-1}\{Y(s)\}$$

$$= \cos(t) + \sin(t)$$

Example

Solve the following integral equation.

$$y(t) + \int_0^t (t-v)y(v) \, dv = t^2$$

$$y(t) + t * y(t) = t^2$$

$$\mathcal{L}\{y(t) + t * y(t)\} = \mathcal{L}\{t^2\}$$

$$Y(s) + \frac{1}{s^2}Y(s) = \frac{2}{s^3}$$

$$\left(1 + \frac{1}{s^2}\right)Y(s) = \frac{2}{s^3}$$

$$\frac{s^2 + 1}{s}Y(s) = \frac{2}{s^3}$$

$$Y(s) = \frac{2}{s^3} \frac{s^2}{s^2 + 1}$$

$$= \frac{2}{s(s^2 + 1)}$$

partial fraction decomposition omitted for brevity

$$= \frac{2}{s} - \frac{2}{s^2 + 1}$$

$$y(t) = \mathcal{L}^{-1}\{Y(s)\}$$


$$= \mathcal{L}^{-1}\left\{\frac{2}{s}\right\} - \mathcal{L}^{-1}\left\{\frac{2}{s^2 + 1}\right\}$$

$$= 2\mathcal{L}^{-1}\left\{\frac{1}{s}\right\} - 2\mathcal{L}^{-1}\left\{\frac{1}{s^2 + 1}\right\}$$

$$= 2 - 2\cos(t)$$

Example

Consider an LRC Series Circuit.



Suppose that the current at time t in the circuit above is governed by the following equation:

$$L \frac{di}{dt} + Ri + \frac{1}{c} + \int_0^t i(v) \, dv = e^t$$

Find the current $i(t)$ when $L = \frac{1}{10}$, $R = 2$, $C = 10$, and

$$E(t) = \begin{cases} 10, & 0 \leq t < 1 \\ 0, & t > 1 \end{cases}$$

with $i(0) = 0$.

$$\begin{aligned}
\frac{1}{10} \frac{di}{dt} + 2i(t) + 10 \int_0^t i(v) dv &= E(t) \\
\frac{1}{10} \frac{di}{dt} + 2i(t) + 10 \int_0^t i(v) dv &= 10 - 10u(t-1) \\
\mathcal{L}\left\{\frac{1}{10} \frac{di}{dt} + 2i(t) + 10 \int_0^t i(v) dv\right\} &= \mathcal{L}\{10 - 10u(t-1)\} \\
\frac{1}{10} sI(s) + 2I(s) + \frac{10}{s} I(s) &= \frac{10}{s} - \frac{10e^{-s}}{s} \\
I(s) \left[\frac{s}{10} + 2 + \frac{10}{s} \right] &= \frac{10}{s} - 10 \frac{e^{-s}}{s} \\
I(s) \left[\frac{(s+10)^2}{10s} \right] &= \frac{10}{s} - 10 \frac{e^{-s}}{s} \\
I(s) &= \frac{10}{s} \frac{10s}{(s+10)^2} - e^{-s} \left[\frac{10}{s} \frac{10s}{(s+10)^2} \right] \\
&= \frac{100}{(s+10)^2} - e^{-s} \frac{100}{(s+10)^2} \\
i(t) &= \mathcal{L}^{-1}\{I(s)\} \\
&= \mathcal{L}^{-1}\left\{\frac{100}{(s+10)^2}\right\} - \mathcal{L}^{-1}\left\{e^{-s} \frac{100}{(s+10)^2}\right\} \\
&= 100te^{-10t} - 100(t-1)e^{-10(t-1)}u(t-1)
\end{aligned}$$

You can find all my notes at <http://omgimanerd.tech/notes>. If you have any questions, comments, or concerns, please contact me at alvin@omgimanerd.tech