

Linear Algebra

Alvin Lin

August 2017 - December 2017

Final Exam Review

1. Solve a linear system
2. Like #8 from Exam 1. Given a 3 by 3 linear system with some coefficients k , find all values of k such that the solution has no solutions, 1 solution, or infinitely many solution.
3. Similar to Exam 2, #3. A square matrix M satisfies a given polynomial equation. Find M^{-1} , $rank(M)$, $nullity(M)$.
4. Given a vector space V and some subsets of V , determine if those subsets are subspaces of V .
5. Like Exam 2, #7. Given a 3 by 3 matrix A , with some entries in terms of k , find all values of k making A invertible.
6. Given 2 vector spaces V , determine if they are isomorphic.
7. Compute some determinants.
8. Similar to Exam 3 #5
9. Similar to Exam 3 #4
10. A question involving linear transformation $T : V \rightarrow W$ and the rank-nullity theorem.
11. Similar to questions from the Exam 4.
12. Similar to questions from the Exam 4.

Bonus: Given some vector space V , compute its dimension.

Bonus: Given transformation $T : V \rightarrow W$, find $[T]_{C \leftarrow B}$ where B is a basis for V and C is a basis for W .

Example

Given a linear transformation $T : V \rightarrow W$ with $\dim(V) = 5$ and $\dim(W) = 6$. Are there any linear transformations $T : V \rightarrow W$ that are also onto? No, the rank-nullity theorem states that $\dim(V) = \text{rank}(T) + \text{nullity}(T)$.

$$\text{rank}(T) = \dim(\text{range}(T)) \leq \dim(V) < \dim(W)$$

$$\text{range}(T) \neq W$$

So T is not onto.

Suppose $\dim(V) = 6$, $\dim(W) = 5$. Are there any one-to-one linear transformations $T : V \rightarrow W$?

$$\text{rank}(T) \leq \dim(W) = 5$$

$$\dim(V) = \text{rank}(T) + \text{nullity}(T)$$

$$\therefore \text{nullity}(T) \geq 1$$

$$\text{nullity}(T) \neq \{\vec{0}\}$$

T is not one-to-one.

Example

V is a vector space with subspaces U, W . Prove $U \cap W$ is a subspace of V .

1. Is $\vec{0} \in U \cap W$? U is a subspace so $\vec{0} \in U$. W is a subspace so $\vec{0} \in W$. Therefore, $\vec{0} \in U \cap W$.
2. Let $\vec{x}, \vec{y} \in U \cap W$. So $\vec{x}, \vec{y} \in U$ and $\vec{x}, \vec{y} \in W$. Then $\vec{x} + \vec{y} \in U$ and $\vec{x} + \vec{y} \in W$. Thus, $\vec{x} + \vec{y} \in U \cap W$.
3. Let $\vec{u} \in U \cap W$ and c be a scalar. $\vec{u} \in U$ and $\vec{u} \in W$. Then $c\vec{u} \in U$ and $c\vec{u} \in W$. Therefore, $c\vec{u} \in U \cap W$.

Example

Let V be a vector space. Take U, W as subspaces of V . Is it true that $U \cap W$ is a subspace of V ? Let $V = \mathbb{R}^2$. $U = \{(x, 0) \mid x \in \mathbb{R}\}$ and $W = \{(0, y) \mid y \in \mathbb{R}\}$. $U \cap W$ is the union of the x and y axes, which is not closed under addition. For $U \cap W$ to be closed under addition, we need $U \subseteq W$ or $W \subseteq U$.

Example

Say B is a set of vectors in a vector space V with the property that every $\vec{v} \in V$ can be written uniquely as a linear combination of elements of B . Explain why B is a basis for V . Show that the $\text{span}(B) = V$.

Take $\vec{v} \in V$

$$\vec{v} = \sum_{i=1}^n c_i \vec{v}_i$$

$$\text{span}(B) = V$$

Show that B is linearly independent.

$$\sum_{i=1}^n c_i \vec{v}_i = \vec{0} = 0\vec{v}_1 + 0\vec{v}_2 + \cdots + 0\vec{v}_n$$

By uniqueness, $c_1 = c_2 = \cdots = c_n = 0$. Thus, B is linearly independent, so B is a basis for V .

You can find all my notes at <http://omgimanerd.tech/notes>. If you have any questions, comments, or concerns, please contact me at alvin@omgimanerd.tech