

University Physics 1A

Alvin Lin

November 2nd, 2017

Angular Motion

A pulley consists of a hub of radius 0.160m attached to a wheel of radius 0.600m. Initially the pulley is rotating clockwise at 6.50 rad/s, and at some time later it is rotating counterclockwise at 2.50 rad/s. The angular position at the second time is 24.0 rad clockwise relative to the first time. A string wrapped around the hub and is attached to a block.

- (a) Find the angular acceleration of the pulley. Be sure to give a vector direction, not using “clockwise” or “counterclockwise”.

$$\begin{aligned}\omega^2 &= \omega_0^2 + 2\alpha\theta \\ (-6.5)^2 &= 2.5^2 + 2\alpha 24.0 \\ \alpha &= 0.75 \frac{\text{rad}}{\text{s}} \quad (+\hat{k} \text{ direction})\end{aligned}$$

- (b) Find the time between the two points mentioned.

$$\begin{aligned}\omega &= \omega_0 + \alpha t \\ 2.5 &= (-6.5) + 0.75t \\ t &= 12\text{s}\end{aligned}$$

- (c) Find the acceleration of the block, including its direction.

$$\begin{aligned}a_{\text{tangential}} &= \alpha r \\ &= 0.75(0.160) \\ &= 0.12 \frac{\text{m}}{\text{s}^2}\end{aligned}$$

- (d) The string from the block touches the pulley at the hub. At the initial time find the centripetal acceleration of that point on the hub.

$$\begin{aligned}a_{centripetal} &= \frac{v^2}{r} \\ &= r\omega^2 \\ &= 0.160(6.5^2) \\ &= 6.76 \frac{m}{s^2}\end{aligned}$$

Moment of Inertia

$$\begin{aligned}KE &= \frac{1}{2}mv^2 \\ &= \frac{1}{2}m(r\omega^2) \\ &= \frac{1}{2}(mr^2)\omega^2\end{aligned}$$

$$I_{ring} = \text{moment of inertia} = mr^2$$

$$\begin{aligned}I_{disk} &= \sum_i m_i r_i^2 \\ &= \int r^2 dm\end{aligned}$$

You can find all my notes at <http://omgimanerd.tech/notes>. If you have any questions, comments, or concerns, please contact me at alvin@omgimanerd.tech