

The Substitution Rule

Alvin Lin

Calculus II: August 2016 - December 2016

The Substitution Rule

$$\int \sin(t)\sqrt{1 + \cos(t)}dt$$

Substitute unsolvable terms with u to reduce the equation to something easier to integrate.

$$\begin{aligned} \text{Let : } \quad u &= 1 + \cos(t) \\ du &= -\sin(t)dt \\ dt &= \frac{-du}{\sin(t)} \end{aligned}$$

Substitute your new terms back into the original equation. The goal is to get a friendlier integral in terms of another variable. If the new integral has both x and u in it, then it may be advisable to try a different method of integration or substituting a different term.

$$\int \sin(t)\sqrt{u}\left(\frac{-du}{\sin(t)}\right)$$

One we simplify the integral after substituting everything, it becomes much easier to integrate.

$$\begin{aligned} & - \int u^{\frac{1}{2}} du \\ & - \left(\frac{u^{\frac{3}{2}}}{\frac{3}{2}}\right) + C \\ & = -\frac{2}{3}(1 + \cos(t))^{\frac{3}{2}} + C \end{aligned}$$

Practice problem 30

$$\int \frac{\sec^2(x)}{\tan^2(x)} dx$$

$$\text{Let : } u = \tan(x)$$

$$du = \sec^2(x) dx$$

$$\int \frac{du}{u^2}$$

$$\int u^{-2} du$$

$$\frac{u^{-1}}{-1} + C$$

$$= -\frac{1}{\tan(x)} + C$$

Practice problem 31

$$\int \frac{\tan^{-1}(x^2)}{1+x^2} dx$$

$$\text{Let : } u = \tan^{-1}(x)$$

$$du = \frac{1}{1+x^2} dx$$

$$\int u^2 du$$

$$= \frac{u^3}{3} + C$$

Practice problem 42

$$\int \frac{\cos(\ln(t))}{t} dt$$

$$\text{Let : } u = \ln(t)$$

$$du = \frac{1}{t} dt$$

$$\int \cos(u) du$$

$$= \sin(u) + C$$

If any errors are found, please contact me at alvin.lin.dev@gmail.com