

Multivariable and Vector Calculus

Alvin Lin

August 2017 - December 2017

Review 1

$$\vec{a} \cdot \vec{b} = a_1b_1 + a_2b_2 + a_3b_3 = |\vec{a}||\vec{b}| \cos \theta$$

$$\text{proj}_{\vec{a}} \vec{b} = \frac{\vec{a} \cdot \vec{b}}{|\vec{a}|} \vec{a}$$

$$\vec{a} \cdot \vec{b} = 0 \equiv \vec{a} \perp \vec{b}$$

$$\vec{a} \times \vec{b} = \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ a_1 & a_2 & a_3 \\ b_1 & b_2 & b_3 \end{vmatrix}$$

$$\vec{a} \times \vec{b} \perp \vec{a}, \vec{b}$$

$$\vec{a} \times \vec{b} = -\vec{b} \times \vec{a}$$

$$\vec{a} \times \vec{b} = \vec{0} \equiv \vec{a} \parallel \vec{b}$$

$$\text{Volume of a pallellepiped} = (\vec{a} \times \vec{b}) \cdot \vec{c} = \begin{vmatrix} c_1 & c_2 & c_3 \\ a_1 & a_2 & a_3 \\ b_1 & b_2 & b_3 \end{vmatrix}$$

$$\text{Line} = \begin{cases} x & = x_o + ta_1 \\ y & = y_o + ta_2 \\ z & = z_o + ta_3 \end{cases}$$

$$\text{Plane} = n_1(x - x_o) + n_2(y - y_o) + n_3(z - z_o) = 0$$

Practice Problem

Find the angle between the diagonal of a cube and the edge of the base starting at $(0,0,0)$.

$$\cos \alpha = \frac{\langle 1, 1, 1 \rangle \cdot \langle 0, 1, 0 \rangle}{|\langle 1, 1, 1 \rangle| |\langle 0, 1, 0 \rangle|} = \frac{1}{\sqrt{3}}$$

Practice Problem

Find x such that $\langle 1, x, 2 \rangle \perp \langle x, 3, 4 \rangle$.

$$\begin{aligned}\langle 1, x, 2 \rangle \cdot \langle x, 3, 4 \rangle &= 0 \\ x + 3x + 8 &= 0 \\ x &= -2\end{aligned}$$

Is there such an x that the two vectors are parallel?

$$\begin{aligned}c\langle 1, x, 2 \rangle &= \langle x, 3, 4 \rangle \\ c &= x \\ cx &= 3 \\ 2c &= 4\end{aligned}$$

This system of linear equations is inconsistent, therefore the two vectors are not parallel.

Practice Problem

Given the planes $\Pi_1 : x - 2y + z = 4$ and $\Pi_2 : 2x - 2x - z = 6$. Find the line of intersection between the planes $l = \Pi_1 \cap \Pi_2$.

If we let $x = 0$:

$$\begin{cases} -2y + z = 4 \\ -2y - z = 6 \end{cases}$$
$$y = -\frac{5}{2} \quad z = -1 \quad P_1 = \left(0, -\frac{5}{2}, 1\right)$$

If we let $y = 0$:

$$\begin{cases} x + z = 4 \\ 2x - z = 6 \end{cases}$$
$$x = \frac{10}{3} \quad z = \frac{2}{3} \quad P_2 = \left(\frac{10}{3}, 0, \frac{2}{3}\right)$$

$$\overrightarrow{P_1P_2} = \left\langle \frac{10}{3}, \frac{5}{2}, -\frac{1}{3} \right\rangle$$

$$l = \begin{cases} x &= \frac{10}{3} + t\frac{10}{3} \\ x &= t\frac{5}{2} \\ z &= \frac{2}{3} - t\frac{1}{3} \end{cases}$$

Find the angle between Π_1, Π_2 .

$$\cos \alpha = \frac{\vec{n}_1 \cdot \vec{n}_2}{|\vec{n}_1||\vec{n}_2|} = \frac{\langle 1, -2, 1 \rangle \cdot \langle 2, -2, -1 \rangle}{\sqrt{6}\sqrt{9}} = \frac{5}{3\sqrt{6}}$$

Practice Problem

Given $\Pi_1 : x - 2y + z = 4$ and $l : x - 2 = y + 5 = z - 4 = t$, find $P = l \cap \Pi$.

$$\begin{aligned} x - 2y + z &= 4 \\ (t + 2) - 2(t - 5) + (t + 4) &= 4 \\ 16 &\neq 4 \end{aligned}$$

Since there is no solution, the line l does not intersect the plane.

Find the distance between l and Π . Pick any point P_0 on the plane and any point P_1 on the line.

$$\begin{aligned} P_0 &= (4, 0, 0) \\ P_1 &= (2, -5, 4) \\ dist(l, \Pi) &= comp_{\vec{n}} \overrightarrow{P_0P_1} \\ &= \frac{\overrightarrow{P_0P_1} \cdot \vec{n}}{|\vec{n}|} \\ &= \frac{\langle -2, -5, 4 \rangle \cdot \langle 1, -2, 1 \rangle}{|\langle 1, -2, 1 \rangle|} \\ &= \frac{14}{\sqrt{6}} \end{aligned}$$

You can find all my notes at <http://omgimanerd.tech/notes>. If you have any questions, comments, or concerns, please contact me at alvin@omgimanerd.tech