

# Linear Algebra

Alvin Lin

August 2017 - December 2017

## Spanning Sets and Linear Independence

We want to know when one vector is a **linear combination** of some other vectors.

$$c_1\vec{v}_1 + c_2\vec{v}_2 + c_3\vec{v}_3$$

For example, is the vector  $\begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix}$  a linear combination of  $\vec{v} = \begin{bmatrix} 1 \\ 0 \\ 3 \end{bmatrix}$  and  $\vec{w} = \begin{bmatrix} -1 \\ 1 \\ -3 \end{bmatrix}$ .

The solution would be to find scalars  $c_1, c_2, c_3$  such that:

$$\begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix} = c_1 \begin{bmatrix} 1 \\ 0 \\ 3 \end{bmatrix} + c_2 \begin{bmatrix} -1 \\ 1 \\ -3 \end{bmatrix}$$

We can treat this as a system of linear equations and solve an augmented matrix:

$$A' = \left[ \begin{array}{cc|c} 1 & -1 & 1 \\ 0 & 1 & 2 \\ 3 & -3 & 3 \end{array} \right]$$

Bring this to reduced row echelon form yields:

$$A' = \left[ \begin{array}{cc|c} 1 & 0 & 3 \\ 0 & 1 & 2 \\ 0 & 0 & 0 \end{array} \right]$$

Thus,  $\begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix}$  is a linear combination of  $\vec{v}, \vec{w}$  through the constants:

$$\begin{bmatrix} c_1 \\ c_2 \end{bmatrix} = \begin{bmatrix} 3 \\ 2 \end{bmatrix}$$

## Describing a Linear System

There are three equivalent ways to specify a linear system:

1.

$$\begin{cases} a_{11}x_1 + a_{12}x_2 + \cdots + a_{1n}x_n & = b_1 \\ \vdots & \\ a_{m1}x_1 + a_{m2}x_2 + \cdots + a_{mn}x_n & = b_m \end{cases}$$

2. Let  $\vec{A}_i$  be a column vector with entries  $a_i$ :

$$x_1\vec{A}_1 + x_2\vec{A}_2 + \cdots + x_n\vec{A}_n = \vec{b}$$

3.  $A\vec{x} = \vec{b}$  where  $A$  is an  $m \times x$  matrix:

$$\vec{x} = \begin{bmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{bmatrix}$$

$$\vec{b} = \begin{bmatrix} b_1 \\ b_2 \\ \vdots \\ b_m \end{bmatrix}$$

where we multiply  $A\vec{x} = \vec{b}$  using:

$$b_i = \text{ith row of } A \times \vec{x}$$

**Theorem:** A system of linear equations with augmented matrix  $[A|\vec{b}]$  is consistent if and only if  $\vec{b}$  is a linear combination of the columns of  $A$ .

## Spanning Sets

Let  $\{\vec{v}_1, \dots, \vec{v}_k\}$  be a set of vectors.

$$\text{Span}(\{\vec{v}_1, \dots, \vec{v}_k\}) = \left\{ \sum_{i=1}^k c_i \vec{v}_i \mid c_i \text{'s are scalars} \right\}$$

The span of:

$$\vec{e}_1 = \begin{bmatrix} 1 \\ 0 \end{bmatrix} \quad \vec{e}_2 = \begin{bmatrix} 0 \\ 1 \end{bmatrix}$$

is equal to:

$$\begin{aligned} \text{Span}(\{\vec{e}_1, \vec{e}_2\}) &= \{c_1\vec{e}_1 + c_2\vec{e}_2 \mid c_1, c_2 \in \mathbb{R}\} \\ &= \left\{ \begin{bmatrix} c_1 \\ 0 \end{bmatrix} + \begin{bmatrix} 0 \\ c_2 \end{bmatrix} \mid c_1, c_2 \in \mathbb{R} \right\} \\ &= \left\{ \begin{bmatrix} c_1 \\ c_2 \end{bmatrix} \mid c_1, c_2 \in \mathbb{R} \right\} \\ &= \mathbb{R}^2 \end{aligned}$$

Is a vector  $\vec{v}$  in  $\text{Span}(\{\vec{u}, \vec{w}\})$ ? Can we find scalars  $c_1, c_2$  such that  $\vec{v} = c_1\vec{u} + c_2\vec{w}$ ? This translates to the following:

$$A = [\vec{u} \quad \vec{w}] \quad \vec{x} = \begin{bmatrix} c_1 \\ c_2 \end{bmatrix}$$

And we are essentially solving a system of linear equations:

$$A' = [A \mid \vec{v}]$$

## Linear Independence

We say vectors  $\vec{v}_1, \vec{v}_2, \dots, \vec{v}_m$  are **linearly independent** if:

$$\sum_{i=1}^n c_i \vec{v}_i = \vec{0} \Rightarrow \forall i (c_i = 0)$$

In terms of matrices, let  $A = [\vec{v}_1 \quad \vec{v}_2 \quad \dots \quad \vec{v}_m]$ . Examining  $A\vec{x} = \vec{0}$ , the only solution for  $\vec{x}$  is:

$$\vec{x} = \begin{bmatrix} c_1 \\ c_2 \\ \dots \\ c_m \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ \dots \\ 0 \end{bmatrix}$$

A set of vectors  $\vec{v}_1, \vec{v}_2, \dots, \vec{v}_m$  is **linearly dependent** if it is not linearly independent.

**Example**

Are the vectors  $\vec{0}, \vec{v}_1, \vec{v}_2, \dots, \vec{v}_m$  linearly dependent or independent?

$$c_1(\vec{0}) + c_2\vec{v}_1 + c_3\vec{v}_2 + \dots + c_{m+1}\vec{v}_m = \vec{0}$$

$c_1$  can be 1, so this set of vectors is linearly dependent.

**Example**

Are the vectors

$$\vec{v} = \begin{bmatrix} 1 \\ 1 \end{bmatrix} \quad \vec{w} = \begin{bmatrix} 1 \\ 2 \end{bmatrix}$$

linearly independent or linearly dependent?

$$c_1\vec{v} + c_2\vec{w} \stackrel{?}{=} \vec{0}$$

$$\begin{bmatrix} 1 & 1 \\ 1 & 2 \end{bmatrix} \begin{bmatrix} c_1 \\ c_2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

This has a unique solution, so  $\vec{v}, \vec{w}$  are linearly independent.

**Example**

The general equation of the plane contains points  $(1,0,3)$ ,  $(-1,1,-3)$ , and the origin. It has equation  $ax + by + cz = 0$ . Find  $a, b, c$ .

$$a(1) + 3c = 0$$

$$a(-1) + 1b - 3c = 0$$

$$\left[ \begin{array}{ccc|c} 1 & 0 & 3 & 0 \\ -1 & 1 & -3 & 0 \end{array} \right] = \left[ \begin{array}{ccc|c} 1 & 0 & 3 & 0 \\ 0 & 1 & 0 & 0 \end{array} \right]$$

$$a + 3c = 0$$

$$a = -3c$$

$$b = 0$$

$$-3cx + cz = 0$$

$$-3x + z = 0$$

### Example

Prove that  $\vec{u}, \vec{v}, \vec{w} \in \text{span}(\vec{u}, \vec{v}, \vec{w})$ .

$$\vec{u} = 1\vec{u} + 0\vec{v} + 0\vec{w}$$

$$\vec{v} = 0\vec{u} + 1\vec{v} + 0\vec{w}$$

$$\vec{w} = 0\vec{u} + 0\vec{v} + 1\vec{w}$$

$$\vec{u}, \vec{v}, \vec{w} \in \text{span}(\vec{u}, \vec{v}, \vec{w})$$

### Example

Prove that  $\vec{u}, \vec{v}, \vec{w} \in \text{span}(\vec{u}, \vec{u} + \vec{v}, \vec{u} + \vec{v} + \vec{w})$ .

$$\vec{u} = 1\vec{u} + 0(\vec{u} + \vec{v}) + 0(\vec{u} + \vec{v} + \vec{w})$$

$$\vec{v} = -1\vec{u} + 1(\vec{u} + \vec{v}) + 0(\vec{u} + \vec{v} + \vec{w})$$

$$\vec{w} = 0\vec{u} - 1(\vec{u} + \vec{v}) + 1(\vec{u} + \vec{v} + \vec{w})$$

$$\vec{u}, \vec{v}, \vec{w} \in \text{span}(\vec{u}, \vec{u} + \vec{v}, \vec{u} + \vec{v} + \vec{w})$$

### Useful Fact

Suppose we have  $m$  vectors in  $\mathbb{R}^n$ , where  $m > n$ . Those vectors are linearly dependent.

$$A = [\vec{v}_1 \quad \vec{v}_2 \quad \dots \quad \vec{v}_m]$$

Examine:  $A\vec{x} = \vec{0}$ . As a consequence of the rank theorem, which states that the number of free variables is equal to the number of columns of  $A$  minus the rank of  $A$ ,  $A\vec{x} = \vec{0}$  has a non-trivial solution, hence the columns of  $A$  are linearly dependent.

### Example

Prove that if  $\vec{u}_1, \dots, \vec{u}_m \in \mathbb{R}^n$  where  $S = \{\vec{u}_1, \dots, \vec{u}_k\}$  and  $T = \{\vec{u}_1, \dots, \vec{u}_k, u_{k+1}, \dots, u_m\}$  that  $\text{span}(S) \subseteq \text{span}(T)$ . Let  $\alpha \in \text{span}(S)$ :

$$\begin{aligned} \alpha &= c_1\vec{u}_1 + c_2\vec{u}_2 + \dots + c_k\vec{u}_k \\ &= c_1\vec{u}_1 + \dots + c_k\vec{u}_k + 0u_{k+1} + 0u_{k+2} + \dots + 0u_m \\ &\in \text{span}(T) \end{aligned}$$

Deduce if  $\mathbb{R}^n = \text{span}(S)$ , then  $\mathbb{R}^n = \text{span}(T)$ .

$$\mathbb{R}^n = \text{span}(S) \subseteq \text{span}(T) \subseteq \mathbb{R}^n$$

$$\text{span}(T) = \mathbb{R}^n$$

### Example

Suppose  $\vec{w}$  is a linear combination of  $\vec{u}_1, \vec{u}_2, \dots, \vec{u}_k$  and each  $\vec{u}_i$  is a linear combination of  $\vec{v}_1, \dots, \vec{v}_m$ . Prove  $\vec{w}$  is a linear combination of  $\vec{v}_1, \dots, \vec{v}_m$ .

$$\begin{aligned}\vec{w} &= \sum_{i=1}^k c_i \vec{u}_i \\ &= \sum_{i=1}^k c_i \left( \sum_{j=1}^m d_{ij} \vec{v}_j \right) \\ &= \sum_{i=1}^k \left( \sum_{j=1}^m c_i d_{ij} \vec{v}_j \right) \\ &= \sum_{j=1}^m \left( \sum_{i=1}^k c_i d_{ij} \right) \vec{v}_j \\ w &\in \text{span}(\vec{v}_1, \dots, \vec{v}_m)\end{aligned}$$

Also suppose each  $\vec{v}_j$  is a linear combination of  $\vec{u}_1, \dots, \vec{u}_k$ . Prove  $\text{span}(\vec{u}_1, \dots, \vec{u}_k) = \text{span}(\vec{v}_1, \dots, \vec{v}_m)$ . Above we proved:

$$\text{span}(\vec{u}_1, \dots, \vec{u}_k) \subseteq \text{span}(\vec{v}_1, \dots, \vec{v}_m)$$

Therefore:

$$\text{span}(\vec{u}_1, \dots, \vec{u}_k) \supseteq \text{span}(\vec{v}_1, \dots, \vec{v}_m)$$

### Example

If the columns of an  $n \times n$  matrix  $A$  are linearly independent as vectors in  $\mathbb{R}^n$ , what is the rank of  $A$ ?

$$\text{rank}(A) = n$$

The columns are linearly independent as  $A\vec{x} = \vec{0}$  has only the trivial solution. So there are no free variables.

### Example

Prove two vectors are linearly dependent if and only if 1 is a scalar multiple of the other.

Say  $\vec{u}$  and  $\vec{v}$  are linearly dependent, there exists scalars  $c_1, c_2$  (not both 0) such that  $c_1\vec{u} + c_2\vec{v} = \vec{0}$ . Without loss of generality, we say that  $c_1 \neq 0$ , then:

$$\begin{aligned}c_1\vec{u} &= -c_2\vec{v} \\ \vec{u} &= \frac{-c_2}{c_1}\vec{v}\end{aligned}$$

Suppose  $\vec{u}$  is a scalar multiple of  $\vec{v}$ .

$$\begin{aligned}\vec{u} &= c\vec{v} \\ c\vec{v} - \vec{u} &= \vec{0}\end{aligned}$$

Thus,  $\vec{u}$  and  $\vec{v}$  are linearly dependent.

You can find all my notes at <http://omgimanerd.tech/notes>. If you have any questions, comments, or concerns, please contact me at [alvin@omgimanerd.tech](mailto:alvin@omgimanerd.tech)