

# Boundary Value Problems: Homework 3

Alvin Lin

August 2018 - December 2018

## Problem 1

Exercises 2.1: problems 4-7

### Exercise 4

Show that the set of functions  $\{\sin(n\pi x), -1 < x < 1, n \in \mathbb{N}\}$  is orthogonal.

$$\begin{aligned} n \in \mathbb{N}, m \in \mathbb{N}, n \neq m \\ (f_n, f_m) &= \int_{-1}^1 \sin(m\pi x) \sin(n\pi x) \, dx \\ &= \int_{-1}^1 \frac{1}{2} (\cos(m\pi x - n\pi x) - \cos(m\pi x + n\pi x)) \, dx \\ &= \frac{1}{2} \left[ \frac{1}{m\pi - n\pi} \sin(m\pi x - n\pi x) + \frac{1}{m\pi + n\pi} \sin(m\pi x + n\pi x) \right]_{-1}^1 \\ &= \frac{1}{2} \left[ \frac{\sin(m\pi - n\pi)}{m\pi - n\pi} + \frac{\sin(m\pi + n\pi)}{m\pi + n\pi} - \frac{\sin(n\pi - m\pi)}{m\pi - n\pi} - \frac{\sin(-m\pi - n\pi)}{m\pi + n\pi} \right] \\ &= \frac{1}{2} (0 + 0 - 0 - 0) \\ &= 0 \end{aligned}$$

Because  $m \in \mathbb{N}$  and  $n \in \mathbb{N}$ , all the sines evaluate to some  $\sin(c\pi)$  where  $c \in \mathbb{N}$ . The sine of any multiple of  $\pi$  is 0.

$$\begin{aligned} n \in \mathbb{N}, m \in \mathbb{N}, n = m \\ (f_n, f_m) &= \int_{-1}^1 \frac{1}{2} (\cos(m\pi x - n\pi x) - \cos(m\pi x + n\pi x)) \, dx \\ &= -\frac{1}{2} \int_{-1}^1 \cos(2n\pi x) \, dx \\ &= -\frac{1}{2} \left[ \frac{\cos(2n\pi x)}{2n\pi} \right]_{-1}^1 \\ &\neq 1 \end{aligned}$$

This set is not orthonormal.

### Exercise 5

Find  $\alpha$  so that  $\{1, x, 1 + \alpha x^2\}$  on  $(-1, 1)$  is orthogonal.

$$\begin{aligned}0 &= (1, 1 + \alpha x^2) \\ &= \int_{-1}^1 1 + \alpha x^2 \, dx \\ &= 2 + \alpha \left[ \frac{x^3}{3} \right]_{-1}^1 \\ &= 2 + \alpha \frac{2}{3} \\ -2 &= \frac{2\alpha}{3} \\ \alpha &= -3\end{aligned}$$

Normalize the set.

$$\begin{aligned}\|1\| &= \sqrt{\int_{-1}^1 1^2 \, dx} \\ &= \sqrt{2} \\ \|x\| &= \sqrt{\int_{-1}^1 x^2 \, dx} \\ &= \sqrt{\frac{2}{3}} \\ \|1 - 3x^2\| &= \sqrt{\int_{-1}^1 (1 - 3x^2)^2 \, dx} \\ &= \sqrt{\int_{-1}^1 1 - 6x^2 + 9x^4 \, dx} \\ &= \sqrt{\left[ x - 2x^3 + \frac{9}{5}x^5 \right]_{-1}^1} \\ &= \sqrt{1 - 2 + \frac{9}{5} + 1 - 2 + \frac{9}{5}} \\ &= \sqrt{\frac{18}{5} - 2} \\ &= \sqrt{\frac{8}{5}}\end{aligned}$$

The normalized set is  $\left\{ \frac{1}{\sqrt{2}}, \frac{x\sqrt{3}}{\sqrt{2}}, \frac{(1-3x^2)\sqrt{5}}{2\sqrt{2}} \right\}$ .

### Exercise 6

Show that the set  $\{1, \cos(\frac{n\pi x}{L}), \sin(\frac{m\pi x}{L})\}, n, m \in \mathbb{N}, -L < x < L$  is orthogonal but not orthonormal.

$$\begin{aligned} (\cos(\frac{n\pi x}{L}), \sin(\frac{m\pi x}{L})) &= \int_{-L}^L \cos(\frac{n\pi x}{L}) \sin(\frac{m\pi x}{L}) \, dx \\ &= \int_{-L}^L \frac{1}{2} (\sin(\frac{n\pi x + m\pi x}{L}) - \sin(\frac{n\pi x - m\pi x}{L})) \, dx \\ &= 0 \\ (1, \sin(\frac{n\pi x}{L})) &= \int_{-L}^L \sin(\frac{n\pi x}{L}) \, dx \\ &= 0 \end{aligned}$$

Because  $\sin$  is an odd function over the symmetric integral  $[-L, L]$ .

$$\begin{aligned} (1, \cos(\frac{n\pi x}{L})) &= \int_{-L}^L \cos(\frac{n\pi x}{L}) \, dx \\ &= \left[ \frac{L}{n\pi} \sin(\frac{n\pi x}{L}) \right]_{-L}^L \\ &= \frac{L}{n\pi} (\sin(n\pi) - \sin(-n\pi)) \\ &= 0 \end{aligned}$$

$(f_m, f_n) = 0, m \neq n$ , therefore the set is orthogonal.

$$\begin{aligned} (\cos(\frac{n\pi x}{L}), \cos(\frac{n\pi x}{L})) &= \sqrt{\int_{-L}^L \cos^2(\frac{n\pi x}{L}) \, dx} \\ &= \sqrt{\int_{-L}^L \frac{1 + \cos(\frac{2n\pi x}{L})}{2}} \\ &= \sqrt{L + \frac{1}{2} \left[ \frac{L}{2n\pi} \sin(\frac{2n\pi x}{L}) \right]_{-L}^L} \\ &\neq 1 \end{aligned}$$

Therefore, the set is not orthonormal.

### Exercise 7

Is the set  $\{\cos(\frac{n\pi x}{2}), n \in \mathbb{N}_0\}, 0 < x < 2$ , orthonormal? If it fails to be orthonormal, write the corresponding orthonormal set.

$$\begin{aligned} n \neq m \\ (\cos(\frac{n\pi x}{2}), \cos(\frac{m\pi x}{2})) &= \int_0^2 \cos(\frac{n\pi x}{2}) \cos(\frac{m\pi x}{2}) \, dx \\ &= \frac{1}{2} \int_0^2 \cos(\frac{n\pi x - m\pi x}{2}) + \cos(\frac{n\pi x + m\pi x}{2}) \\ &= \left[ \sin(n\pi x - m\pi x) + \sin(n\pi x + m\pi x) \right]_0^2 \\ &= \sin(2n\pi - 2m\pi) + \sin(2n\pi + 2m\pi) - \sin(0) - \sin(0) \\ &= 0 \end{aligned}$$

The set is orthogonal.

$$\begin{aligned} n = m \\ (\cos(\frac{n\pi x}{2}), \cos(\frac{n\pi x}{2})) &= \sqrt{\int_0^2 \cos^2(\frac{n\pi x}{2}) dx} \\ &= \sqrt{\frac{1}{2} \int_0^2 1 + \cos(n\pi x) dx} \\ &= \sqrt{\frac{1}{2} \left[ x + \frac{\sin(n\pi x)}{n\pi} \right]_0^2} \\ &= \sqrt{\frac{1}{2} [2 + 0 - 0 - 0]} \\ &= \sqrt{1} \\ &= 1 \end{aligned}$$

The set is also orthonormal.

Exercises 3.2: problems 6-8

### Exercise 6

Prove that the sum of two odd functions is odd.

$$\begin{aligned} f(-x) &= -f(x) \\ \int_{-L}^L f(x) dx &= 0 \\ g(-x) &= -g(x) \\ \int_{-L}^L g(x) dx &= 0 \\ \int_{-L}^L f(x) + g(x) dx &= \int_{-L}^L f(x) dx + \int_{-L}^L g(x) dx \\ &= 0 + 0 \\ &= 0 \end{aligned}$$

### Exercise 7

Show that if  $f$  is odd, then  $|f|$  and  $f^2$  are even functions.

$$\begin{aligned} f(-x) &= -f(x) \\ |f(-x)| &= |-f(x)| \\ f(-x) &= f(x) \end{aligned}$$

An odd function is symmetric about the origin. Taking the absolute value reflects the negative portion of the function across the x axis and because of the original symmetry, the function is now symmetric about

the y axis.

$$\begin{aligned}f(-x) &= -f(x) \\f(-x)^2 &= (-f(x))^2 \\f(-x)^2 &= f(x)^2\end{aligned}$$

Squaring an odd function takes the negative portion of the function and reflects it across the x axis due to the change in sign while increasing the whole function by a factor of itself. Because of the original symmetry, this function is now symmetric about the y axis.

## Problem 2

Classify the following functions as even, odd, or neither. Show steps to justify your answers.

(a)  $f(x) = (1 - x^2)^{-\frac{1}{2}}$

$$\begin{aligned}f(a) &= f(-a) \\(1 - a^2)^{-\frac{1}{2}} &= (1 - a^2)^{-\frac{1}{2}}\end{aligned}$$

Even.

(b)  $f(x) = e^{-x} \cos(3x)$

$$\begin{aligned}f(a) &= f(-a) \\e^{-a} \cos(3a) &= e^a \cos(-3a) \\e^{-a} \cos(3a) &\neq e^a \cos(3a)\end{aligned}$$

Neither.

(c)  $f(x) = \sinh(x)$

$$\begin{aligned}f(a) &= f(-a) \\\sinh(a) &= \sinh(-a) \\\sinh(a) &\neq -\sinh(a)\end{aligned}$$

Odd.

If you have any questions, comments, or concerns, please contact me at [alvin@omgimanerd.tech](mailto:alvin@omgimanerd.tech)