

# University Physics 1A

Alvin Lin

December 6th, 2017

## Waves

$$v = \frac{\omega}{k} = f\lambda$$

$$\text{speed of wave} = v = \sqrt{\frac{\text{elastic property}}{\text{inertial property}}}$$

$$\text{for a string(spring)} \quad v = \sqrt{\frac{\text{tension}}{\text{mass density}}} = \sqrt{\frac{F_{\text{tension}}}{\lambda}}$$

$$\text{for air(sound)} \quad v = \sqrt{\frac{\text{bulk modulus of air}}{\text{volume mass density}}} = \sqrt{\frac{B}{\rho}}$$

Principle of superposition: two waves at the same point just add their values. While this does not hold for other concepts, it holds for most of the mediums that we will work with.

## Interference

Two waves going in the same direction with the same amplitude:

$$\begin{aligned} y &= y_{\max} \sin(kx - \omega t) + y_{\max} \sin(kx - \omega t + \phi) \\ &= 2y_{\max} \cos\left(\frac{\phi}{2}\right) \sin\left(kx - \omega t + \frac{\phi}{2}\right) \end{aligned}$$

If  $\phi = \pi$  and the two waves are half a wavelength apart, then the two waves are destructive and cancel each other out. If  $\phi = 0$ , then the two are constructive and

amplify each other. Any other values of  $\phi$  are partially constructive. This is an example of wave interference.

Two waves travelling in the opposite direction:

$$\begin{aligned}y &= y_{max} \sin(kx - \omega t) - y_{max} \sin(kx + \omega t) \\ &= 2y_{max} \cos(\omega t) \sin(kx)\end{aligned}$$

This forms a standing wave.

You can find all my notes at <http://omgimanerd.tech/notes>. If you have any questions, comments, or concerns, please contact me at [alvin@omgimanerd.tech](mailto:alvin@omgimanerd.tech)